



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Simply Transitive Primitive Groups Whose Maximal Subgroup Contains a Transitive Constituent of Order p^2 , or pq , or a Transitive Constituent of Degree 5.

BY ELIZABETH RUTH BENNETT.

If the maximal subgroup G_1 of degree $n-1$ of a simply transitive primitive group G of degree n contains a transitive constituent of degree 3, it is known that it must also contain another transitive constituent of degree 3 or 6.* Similar restrictions on G_1 have been determined when G_1 contains a transitive constituent of degree 4.† We proceed to consider restrictions that may be placed on the degree of G or on the transitive constituents of G_1 when G_1 contains a transitive constituent of order p^2 or pq , or a transitive constituent of degree 5.

THEOREM I. *If a transitive constituent of G_1 is of order p^2 , the degree of G is q^a , q a prime. When k , the number of transitive constituents of order p^2 in G_1 , is odd, $k \equiv 3, \text{ mod. } 4$, and $q=2$. When $k \equiv 2, \text{ mod. } 4$, q is a prime of form $4n+3$ and a is an odd number.*

All groups of order p^2 are abelian and can be represented transitively only in regular form. Therefore the order of G_1 can not exceed p^2 and G_1 must be formed from the simple isomorphism of groups of degree and order p^2 .‡ G is then of class $n-1$ and of degree q^a , q a prime.§

The following relation must exist where k represents the number of transitive constituents of order p^2 in G_1 :

$$kp^2 + 1 = q^a. \tag{I}$$

By considering (I) with respect to modulus 4, the remainder of the theorem is evident.

THEOREM II. *If a transitive constituent of G_1 is of order pq , p and q primes, $p > q$, and the order of G_1 is pq , G_1 is formed from establishing a*

* Miller, AMERICAN JOURNAL OF MATHEMATICS, Vol. XXXV, p. 7.

† Bennett, AMERICAN JOURNAL OF MATHEMATICS, Vol. XXXIV, pp. 8, 9.

‡ Miller, *Bulletin American Mathematical Society*, Vol. VI, p. 104.

§ Frobenius, *Berliner Sitzungsberichte* (1902), pp. 455-459.

simple isomorphism between groups of order pq . When the constituent of order pq is abelian, G is of degree r^a , r a prime, and of class $n-1$. If the order of G_1 exceeds pq , G_1 must contain a transitive constituent of degree pq whose order is greater than pq , and, in case the constituent of order pq is of degree p , the order of G_1 must be $q^a p$.

When the order of G_1 is pq , G_1 is formed from the simple isomorphism of groups of order pq , for the order of G_1 is not divisible by the square of a prime number.*

A group of order pq can be represented transitively only on pq or p letters, and, in case the group of order pq is abelian, the representation must be on pq letters. Therefore, if the order of G_1 is pq and the constituent of order pq is abelian, G is of class $n-1$ and degree r^a , r a prime. When the order of G_1 exceeds pq and the transitive constituent of order pq is regular, then G_1 must contain an additional transitive constituent of degree pq whose order exceeds pq .† When the constituent of order pq is of degree p , the subgroup leaving a letter of the constituent of degree p fixed is composed of transitive constituents of order and degree q . G_1 must then contain a transitive constituent whose degree is pq and whose order is greater than pq .‡ If the constituent of order pq is of degree p , the order of G_1 must be $q^a p$,† for the order of G_1 is not divisible by p^2 .

COROLLARY I. If G_1 contains a dihedral group of prime degree p as a transitive constituent, the order of G_1 is $2^a p$.

THEOREM III. When G_1 contains k transitive constituents of order 5, k an odd number, the degree of G is $2^{4\beta}$ and $k \equiv 3, \text{ mod. } 4$. If $k \equiv 2, \text{ mod. } 4$, the degree of G is q^a , q a prime of form $4n+3$ and a an odd number.

Since G_1 contains a transitive constituent of order 5, the order of G_1 is 5 and G is of class $n-1$. The degree of G is then q^a , q a prime. If the number of transitive constituents of order 5 is odd, $5k+1$ is even or $5k+1=2^a$ and $k \equiv 3, \text{ mod. } 4$. Since 2 is a primitive root of 5, in order that the above equation be satisfied $a=4\beta$. If $k \equiv 2, \text{ mod. } 4$, $5k+1 \equiv 3, \text{ mod. } 4$, and in order that $q^a \equiv 3, \text{ mod. } 4$, a must be odd and q a prime of form $4n+3$.

THEOREM IV. If G_1 contains the semi-metacyclic group of degree 5 as a transitive constituent, G_1 must contain another transitive constituent of degree 10 or 5 and the order of G_1 must be $2^a \cdot 5$.

* Miller, *Proceedings London Mathematical Society*, Vol. XXVIII, p. 536.

† Reitz, *AMERICAN JOURNAL OF MATHEMATICS*, Vol. XXVI, p. 9.

‡ Bennett, *loc. cit.*, p. 6.

When the order of G_1 is 10, G_1 can be formed only from the simple isomorphism of groups of degrees 5 and 10. If the order of G_1 exceeds 10, from Theorem II G_1 must contain a transitive constituent of degree 10, and the order of G_1 is $2^a \cdot 5$.

THEOREM V. *If G_1 contains the alternating group of degree 5 as a transitive constituent and the order of G_1 is 60, G_1 is formed from the simple isomorphism of groups whose degree can be only 60, 30, 20, 15, 12, 10, 6 and 5. If the order of G_1 exceeds 60, G_1 must contain a transitive constituent of degree 20. The order of G_1 is $2^a \cdot 3^b \cdot 5$.*

When the order of G_1 is 60, since the alternating group of degree 5 in simple, the order of the other transitive constituents of G_1 must also be 60. The group of order 60 can be represented only on 60, 30, 20, 15, 12, 10, 6 and 5 letters; therefore, only transitive constituents of such degrees may occur when G_1 is of order 60. If the order of G_1 exceeds 60, G_1 must contain a transitive constituent of degree 20, for the subgroup leaving fixed a letter of the alternating group of degree 5 is a primitive group.* Since the order of G_1 is not divisible by 5^2 , the order of G is $2^a \cdot 3^b \cdot 5$.

A theorem concerning the symmetric group of degree 5 may be stated which differs from Theorem V only with regard to the possible representations.

* Bennett, *loc. cit.*, p. 6.